arXiv:0804.3938v1 [math.PR] 24 Apr 2008

Quantum Gross Laplacian and Applications

Samah Horrigue and **Habib Ouerdiane**
Department of Mathematics
Faculty of sciences of Tunis
University of Tunis El-Manar
1060 Tunis, Tunisia
e-mail: samah.horrigue@fst.rnu.tn
e-mail: habib.ouerdiane@fst.rnu.tn

April 24, 2008

Abstract

In this paper, we introduce and study a noncommutative extension of the Gross Laplacian, called quantum Gross Laplacian. Then, applying the quantum Gross Laplacian to the particular case where the operator is the multiplication operator, we find a relation between classical and quantum Gross Laplacian. As application, we give explicit solution of linear quantum white noise differential equation. In particular, we give a explicit solution of the quantum Gross heat equation.

Keywords and phrases. Space of entire functions with growth condition, quantum Gross Laplacian, symbols of operators, convolution operators, quantum Gross heat equations.
Mathematics Subject Classification: 60H40, 60H15, 46F25, 46G20, 81S25.

1 Introduction

The Gross Laplacian Δ_G was introduced by L. Gross in [6] in order to study differential equations in infinite dimensional spaces. It has been shown that the solution of the Cauchy problem

$$\frac{\partial}{\partial t}U(t) = \frac{1}{2}\Delta_G U(t), \quad U(0) = \varphi \tag{1.1}$$

is represented as an integral with respect to Gaussian measure, see [6] and [15]. There exists many literature dedicated to the Gross Laplacian with different points of view. We would like to mention the white noise analysis approach, see [2, 7, 8, 10] and references therein. In [4] and [1], using the fact that the Gross Laplacian is a convolution operator, the authors applied Laplace transform techniques to solve the Cauchy problem (1.1). Moreover, for a smooth initial condition the solution is represented as an integral with respect to a Gaussian measure.

The main purposes of this work are the following: one is to define and study the generalized Gross Laplacian acting on operators denoted Δ_G^Q. Another one is to solve linear quantum stochastic differential equations. In particular, we give explicit solutions of the quantum Gross heat equations.

The paper is organized as follows. In section 2, we review from [9] basic concepts, definitions and results essential to know the space of test and generalized functions denoted respectively $\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$ and $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$. In section 3, we introduce and define the generalized Gross Laplacian Δ_G on the space of entire functions with exponential growth of finite type in two infinite dimensional variables. Then, we prove in Theorem 3.2, that the Gross Laplacian is a convolution operator on the test functions space $\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$, i. e., there exists a distribution $\mathcal{T} \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$, such that

$$\Delta_G \varphi = \mathcal{T} * \varphi,\ \varphi \in \mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2'). \tag{1.2}$$

The relation (1.2) permit to us to extend in natural way the Gross Laplacian applied to the distributions spaces $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$ as follows

$$\Delta_G \Phi = \mathcal{T} * \Phi,\ \Phi \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2'). \tag{1.3}$$

Using the Schwartz-Grothendieck Kernel Theorem and the definition (1.3), we introduce and study, in section 4, the noncommutative extension of Gross Laplacian, denoted by Δ_G^Q, acting on $\mathcal{L}(\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2'), \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2'))$ the space of continuous linear operators from $\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$ into $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$. Then, we establish an analytic characterization of the quantum Gross Laplacian (see Proposition 3.3), i. e., for all $\Xi \in \mathcal{L}(\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2'), \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2'))$, we have

$$\sigma(\Xi)(\xi_1,\xi_2) = (\langle \xi_1, \xi_1 \rangle_1 + \langle \xi_2, \xi_2 \rangle_2)\, \sigma(\Xi)(\xi_1,\xi_2),\ (\xi_1,\xi_2) \in N_1 \times N_2, \tag{1.4}$$

where $\sigma(\Xi)$ denoted the symbol of the operator Ξ. In section 5, we study the action of the quantum Gross Laplacian to the multiplication operator $\mathcal{M}_\Phi$ defined in (5.1) where $\Phi \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$, and we prove that:

$$\left(\Delta_G^Q \mathcal{M}_\Phi \right) e_0 = \Delta_G \Phi, \tag{1.5}$$

where e_0 is the vacuum vector. Therefore, the equality (1.5) establish a relation between the classical and quantum Gross Laplacian. In section 6, we give in Theorem 6.2, the solution of the following quantum stochastic differential equation

$$(E) \left\{ \begin{array}{l} \frac{d\Xi(t)}{dt} = Z(t) * \Xi(t) + \Theta(t), \\ \Xi(0) = \Xi_0, \end{array} \right. \tag{1.6}$$

where $t \mapsto Z(t)$ and $t \mapsto \Theta(t)$ are continuous operator valued process defined on an interval containing the origin $I \subset \mathbb{R}$, i. e.,

$$\begin{array}{rcl} Z : t \in I & \rightarrow & Z(t) \in \mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2')), \\ \Theta : t \in I & \rightarrow & \Theta(t) \in \mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2')) \end{array}$$

and the initial condition $\Xi_0 \in \mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2'))$. As an application of Theorem 6.2, we give explicit solutions of the heat equation associated with the quantum Gross Laplacian.

2 Preliminaries

For $i = 1, 2$, let N_i be a complex nuclear Fréchet space whose topology is defined by a family of increasing Hilbertian norms $\{|.|_{i,p}, p \in \mathbb{N}\}$. For $p \in \mathbb{N}$, we denote by $(N_i)_p$ the completion of N_i with respect to the norm $|.|_{i,p}$ and by $(N_i)_{i,-p}$ respectively N_i' the strong dual space of $(N)_p$ and N. Then, we obtain

$$N_i = \operatorname{proj}\lim_{p\to\infty} (N_i)_p \text{ and } N_i' = \operatorname{ind}\lim_{p\to\infty} (N_i)_{-p}. \tag{2.1}$$

The spaces N_i and N_i' are respectively equipped with the projective and inductive limit topology. For all $p \in \mathbb{N}$, we denote by $|.|_{i,-p}$ the norm on $(N_i)_{-p}$ and by $\langle .,. \rangle_i$ the $\mathbb{C}$-bilinear form on $N_i' \times N_i$.

In the following, H denote the direct Hilbertian sum of $(N_1)_0$ and $(N_2)_0$, i. e., $H = (N_1)_0 \oplus (N_2)_0$.

For $n \in \mathbb{N}$, we denote by $N_i^{\widehat{\otimes} n}$ the $n-$fold symmetric tensor product on N_i equipped with the $\pi-$topology and by $(N_i)_p^{\widehat{\otimes} n}$ the $n-$fold symmetric Hilbertian tensor product on $(N_i)_p$. We will preserve the notation $|.|_{i,p}$ and $|.|_{i,-p}$ for the norms on $(N_i)_p^{\widehat{\otimes} n}$ and $(N_i)_{-p}^{\widehat{\otimes} n}$, respectively.

Let θ be a Young function, i. e., it is a continuous, convex and increasing function defined on $\mathbb{R}^+$ and satisfies the two conditions: $\theta(0) = 0$ and $\lim_{x\to+\infty} \frac{\theta(x)}{x} = +\infty$. Obviously, the conjugate function θ^* of θ defined by

$$\forall x \geq 0, \quad \theta^*(x) := \sup_{t\geq 0}(tx - \theta(t)), \tag{2.2}$$

is also a Young function. For every $n \in \mathbb{N}$, let

$$\theta_n = \inf_{r>0} \frac{e^{\theta(r)}}{r^n}. \tag{2.3}$$

2.1 Spaces of entire functions with growth condition

Throughout the paper, we fix a pair of Young functions (θ_1, θ_2). For all pair of positive numbers $a_1, a_2 > 0$ and pair of integers $(p, q) \in \mathbb{N} \times \mathbb{N}$, we define the space of all entire functions on $(N_1)_{-p} \times (N_2)_{-q}$ with $(\theta_1, \theta_2)-$exponential growth by

$$Exp((N_1)_{-p}\times(N_2)_{-q}, (\theta_1, \theta_2), (a_1, a_2)) = \{f \in \mathcal{H}(N_1\times N_2); \|f\|_{(\theta_1,\theta_2),(a_1,a_2)} < \infty\},$$

where $\mathcal{H}(N_1 \times N_2)$ is the space of all entire functions on $N_1 \times N_2$ and

$$\|f\|_{(\theta_1,\theta_2),(a_1,a_2)} = \sup\{|f(z_1, z_2)|e^{-\theta_1(a_1|z_1|_{-p})-\theta_2(a_2|z_2|_{-q})}, (z_1, z_2) \in (N_1)_{-p}\times(N_2)_{-q}\}.$$

So, the space of all entire functions on $(N_1)_{-p}\times(N_2)_{-q}$ with $(\theta_1, \theta_2)-$exponential growth of minimal type is naturally defined by

$$\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2') = \operatorname{proj}\lim_{\substack{p,q\to\infty \\ a_1,a_2\downarrow 0}} Exp((N_1)_{-p} \times (N_2)_{-q}, (\theta_1, \theta_2), (a_1, a_2)). \tag{2.4}$$

Similarly, the space of entire functions on $N_1 \times N_2$ with $(\theta_1, \theta_2)-$exponential growth of finite type is defined by

$$\mathcal{G}_{(\theta_1,\theta_2)}(N_1 \times N_2) = \operatorname{ind}\lim_{\substack{p,q\to\infty \\ a_1,a_2\to 0}} Exp((N_1)_p \times (N_2)_q, (\theta_1, \theta_2), (a_1, a_2)). \tag{2.5}$$

By definition, $\varphi \in \mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$ and $\Psi \in \mathcal{G}_{(\theta_1,\theta_2)}(N_1 \times N_2)$ admit the Taylor expansions:

$$\varphi(x,y) = \sum_{n,m\in\mathbb{N}} \langle x^{\otimes n} \otimes y^{\otimes m}, \varphi_{n,m}\rangle,\ (x,y) \in N_1' \times N_2', \tag{2.6}$$

and

$$\Psi(\xi,\eta) = \sum_{n,m\in\mathbb{N}} \langle \Psi_{n,m}, \xi^{\otimes n} \otimes \eta^{\otimes m}\rangle,\ (\xi,\eta) \in N_1 \times N_2, \tag{2.7}$$

where for all $n,m \in \mathbb{N}$, we have $\varphi_{n,m} \in N_1^{\hat{\otimes} n} \otimes N_2^{\hat{\otimes} m}$, $\Psi_{n,m} \in \left(N_1^{\hat{\otimes} n}\right)' \otimes \left(N_2^{\hat{\otimes} m}\right)'$ and we used the common symbol $\langle\, .\, ,\, .\, \rangle$ for the canonical $\mathbb{C}-$bilinear form on $\left(N_1^{\otimes n} \times N_2^{\otimes m}\right)' \times N_1^{\otimes n} \times N_2^{\otimes m}$. So, we identify in the next all test function $\varphi \in \mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$ (resp. all generalized function $\Psi \in \mathcal{G}_{(\theta_1,\theta_2)}(N_1 \times N_2)$) by their coefficients of its Taylor series expansion at the origin $(\varphi_{n,m})_{n,m\in\mathbb{N}}$ (resp. $(\Psi_{n,m})_{n,m\in\mathbb{N}}$).

Denote by $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$ the topological dual of $\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$, called the space of distributions on $N_1' \times N_2'$.

For a fixed $(\xi,\eta) \in N_1 \times N_2$, the exponential function $e_{(\xi,\eta)} \in \mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$ is defined by

$$e_{(\xi,\eta)}(z,t) = \exp\{\langle z,\xi\rangle_1 + \langle t,\eta\rangle_2\},\ (z,t) \in N_1' \times N_2'.$$

Then for every $\Phi \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$, the Laplace transform L of Φ is defined by

$$L\Phi(\xi,\eta) = \widehat{\Phi}(\xi,\eta) = \langle\langle \Phi, e_{(\xi,\eta)}\rangle\rangle. \tag{2.8}$$

Theorem 2.1 *[9] For $i = 1,2$, let N_i be complex nuclear Fréchet space and θ_i a Young function. Then, the Laplace transform L is a topological isomorphism:*

$$L : \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2') \to \mathcal{G}_{(\theta_1,\theta_2)}(N_1 \times N_2). \tag{2.9}$$

Remark 2.1 *In the particular case where $\theta_1 = \theta_2 = \theta$, $N_1 = N$ and $N_2 = \{0\}$. We obtain the following identification have*

$$\mathcal{F}_{(\theta,\theta)}(N' \times \{0\}) = \mathcal{F}_\theta(N')$$

and therefore

$$\mathcal{F}^*_{(\theta,\theta)}(N' \times \{0\}) = \mathcal{F}^*_\theta(N').$$

So the space $\mathcal{F}_{(\theta_1,\theta_2)}(N' \times N_2')$ can be considered as a generalization of the space $\mathcal{F}_\theta(N')$ studied in [5].

2.2 Convolution operators

Let $(z,t) \in N_1' \times N_2'$, the translation operator denoted $\tau_{(z,t)}$ is a linear continuous operator $\tau_{(z,t)}$ from $\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$ into itself is defined by

$$\tau_{(z,t)}\varphi(x,y) = \varphi(x+z, y+t),\ (x,y) \in N_1' \times N_2'. \tag{2.10}$$

The convolution product of the distribution $\Phi \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ with a test function $\varphi \in \mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ is defined as follows

$$\Phi * \varphi(z,t) = \langle\langle \Phi, \tau_{(z,t)}\varphi \rangle\rangle,\ (z,t) \in N'_1 \times N'_2. \tag{2.11}$$

Hence, the convolution product of two distribution $\Psi_1, \Psi_2 \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ is given by

$$\widehat{\Psi_1 * \Psi_2} = \widehat{\Psi_1}\widehat{\Psi_2},\ \Psi_1, \Psi_2 \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N'_1 \times N'_2). \tag{2.12}$$

So, by formula (2.12), the convolution product of distribution is commutative and associative.

We denote by $\mathcal{L}(\mathcal{F}_{\theta_1}(N'_1), \mathcal{F}^*_{\theta_2}(N'_2))$ the space of continuous linear operators from $\mathcal{F}_{\theta_1}(N'_1)$ into $\mathcal{F}^*_{\theta_2}(N'_2)$ endowed with the bounded convergence topology.

A convolution operator on the test space $\mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ is defined as a continuous linear operator from $\mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ into itself which commutes with translation operators. Then, T is a convolution operator on $\mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ if and only if there exists a distribution $\Phi_T \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ such that

$$T(\varphi) = T_\Phi(\varphi) = \Phi_T * \varphi,\ \varphi \in \mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2).$$

Note that the convolution product of two distributions $\Psi_1, \Psi_2 \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ is also given by

$$\langle\langle \Psi_1 * \Psi_2, \varphi \rangle\rangle = \langle\langle \Psi_1, \Psi_2 * \varphi \rangle\rangle,\ \varphi \in \mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2).$$

It follows that the notion of convolution operator T_Φ can be extended to the distribution space $\mathcal{F}^*_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ as follows: $\mathbf{T}$ is a convolution operator on $\mathcal{F}^*_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ if and only if there exist $\Phi_T \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ such that

$$\mathbf{T}(\Psi) = \mathbf{T}_\Phi(\Psi) = \Phi_T * \Psi,\ \Psi \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N'_1 \times N'_2).$$

We remark that this extension of convolution operator $\mathbf{T}$ coincide with the adjoint of T_Φ denoted T^*_Φ. In fact, for all $\Psi \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ and $\varphi \in \mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$, we have:

$$\begin{aligned} \langle\langle T^*_\Phi \Psi, \varphi \rangle\rangle &=: \langle\langle \Psi, T_\Phi \varphi \rangle\rangle \\ &= \langle\langle \Psi, \Phi * \varphi \rangle\rangle \\ &= \langle\langle \mathbf{T}_\Phi \Psi, \varphi \rangle\rangle. \end{aligned}$$

Let $i = 1, 2$. Recall now the notion of the right contraction of order $k \in \mathbb{N}$:

For $n, m \in \mathbb{N} - \{0\}$ and $0 \le k \le m \wedge n$, we denote by $\langle .\,, . \rangle_{i,(k)}$ the bilinear map from $N_i'^{\hat{\otimes} m} \times N_i^{\hat{\otimes} n}$ into $N_i'^{\hat{\otimes} m-k} \hat{\otimes} N_i^{\hat{\otimes} n-k}$ defined by

$$\langle x^{\otimes m}, y^{\otimes n} \rangle_{i,(k)} := \langle x, y \rangle_i^k x^{\otimes(m-k)} \otimes y^{\otimes(n-k)},\ x \in N'_i, y \in N_i.$$

The bilinear map $\langle .\,, . \rangle_{i,(k)}$ is continuous. Then using the density of the vector space generated by $\{x^{\otimes m},\ x \in N'_i\}$ in $N_i'^{\hat{\otimes} m}$ and the vector space generated by $\{y^{\otimes n},\ y \in N_i\}$ in $N_i^{\hat{\otimes} n}$, we can extend $\langle .\,, . \rangle_{i,(k)}$ to $N_i'^{\hat{\otimes} m} \times N_i^{\hat{\otimes} n}$. For all $\Phi_n \in N_i'^{\hat{\otimes} m}$ and $\varphi_n \in N_i^{\hat{\otimes} n}$, $\langle \Phi_n, \varphi_n \rangle_{i,(k)}$ is called the right contraction of Φ_n and φ_n of order k.

It follows that for $(\varphi_1,\varphi_2)\in N_1^{\widehat{\otimes}(n+i)}\times N_2^{\widehat{\otimes}(m+j)}$ and $(\psi_1,\psi_2)\in N_1^{\widehat{\otimes}(n+k)}\times N_2^{\widehat{\otimes}(m+l)}$, we define the following generalized contraction as follows:

$$\langle \varphi_1\otimes\varphi_2,\psi_1\otimes\psi_2\rangle_{n,m}=\langle\varphi_1,\psi_1\rangle_{1,(n)}\langle\varphi_2,\psi_2\rangle_{2,(m)}.$$

So, for $i=1,2$, $\Phi_i\in(N_i')^{\widehat{\otimes}(p_i+m_i)}$, $\varphi_i\in N_i^{\widehat{\otimes}(p_i+n_i)}$ and $\psi_i\in N_i^{\widehat{\otimes}(m_i+n_i)}$, it holds:

$$\langle\langle\Phi_1\otimes\Phi_2,\varphi_1\otimes\varphi_2\rangle_{p_1,p_2},\psi_1\otimes\psi_2\rangle=\langle\Phi_1\otimes\Phi_2,\langle\varphi_1\otimes\varphi_2,\psi_1\otimes\psi_2\rangle_{n_1,n_2}\rangle.$$

By an easy calculation like in [3] , we obtain the following lemma.

Lemma 2.2 *For all test function $\varphi=(\varphi_{n,m})_{n,m\in\mathbb{N}}\in\mathcal{F}_{(\theta_1,\theta_2)}(N_1'\times N_2')$ and generalized function $\Phi=(\Phi_{n,m})_{n,m\in\mathbb{N}}\in\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1'\times N_2')$, we have*

$$\begin{aligned}&\Phi*\varphi(x,y)\\&=\sum_{k,l\in\mathbb{N}}\frac{(n+k)!}{k!}\frac{(m+l)!}{l!}\left\langle x^{\otimes k}\otimes y^{\otimes l},\langle\sum_{n,m\in\mathbb{N}}\Phi_{n,m},\varphi_{n+k,m+l}\rangle_{n,m}\right\rangle,\end{aligned}\tag{2.13}$$

for all $(x,y)\in N_1'\times N_2'$.

3 Generalized Gross Laplacian

Let $F\in C^2(N)$. Then for each $\xi\in N$ there exist $F'(\xi)\in N'$ and $F''(\xi)\in(N\otimes N)'$ such that

$$F(\xi+\eta)=F(\xi)+\langle F'(\xi),\eta\rangle+\frac{1}{2}\langle F''(\xi),\eta\otimes\eta\rangle+o(|\eta|_p^2),\ \eta\in N,\tag{3.1}$$

for some $p\in\mathbb{N}$. Moreover, both maps $\xi\mapsto F'(\xi)\in N'$ and $\xi\mapsto F''(\xi)\in(N\otimes N)'$ are continuous. For notation simplicity, taking into account the canonical isomorphism $(N\otimes N)'\simeq\mathcal{L}(N,N')$, which follows from the kernel theorem for a nuclear space, we write $\langle F''(\xi),\eta\otimes\eta\rangle=\langle F''(\xi)\eta,\eta\rangle=F''(\xi)(\eta,\eta)$.

Let τ be the trace operator defined on $N^{\otimes 2}$ by the formula

$$\langle\tau,\xi\otimes\eta\rangle=\langle\xi,\eta\rangle,\forall\xi,\eta\in N.\tag{3.2}$$

Using the definition of the Gross Laplacian Δ_G given in [6], the authors studied in [3], the action of Δ_G on $\mathcal{F}_\theta(N')$. Then, they prove that Δ_G is in fact a convolution operator i. e., for all $\varphi\in\mathcal{F}_\theta(N')$, we have

$$\begin{aligned}\Delta_G\varphi(x)&:=trace_H D^2\varphi(x)\\&=\sum_{n\in\mathbb{N}}(n+2)(n+1)\langle x^{\otimes^n},\langle\tau,\varphi_{n+2}\rangle\rangle\qquad(3.3)\\&=\mathcal{T}*\varphi(x),\ x\in N',\qquad(3.4)\end{aligned}$$

where the Taylor expansion of the distribution $\mathcal{T}$ is given by

$$\mathcal{T}_n=\begin{cases}0,\ n\neq 2\\ \tau,\ n=2.\end{cases}\tag{3.5}$$

Then in the paper [1] , the authors extended the action of the Gross Laplacian on the test function space given in (3.4) to the distribution space as follows

$$\Delta_G \Phi = \mathcal{T} * \Phi,\ \Phi \in \mathcal{F}^*_\theta(N'). \tag{3.6}$$

In this section, we define first the action of the Gross Laplacian on the test function space of two infinite dimensional variables $\mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$.

For $i = 1, 2$, let $\{e^i_j\}_{j\in\mathbb{N}}$ be a complete orthonormal basis of $(N_i)_0$ such that $e^i_j \in N_i$. In all the remainder of this part, we denote by τ_i the trace operator on $N_i^{\otimes 2}$ defined in (3.2).

Lemma 3.1 *[12] For all $i = 1, 2$ and $w_i \in N_i^{\otimes 2}$, it holds that*

$$\langle \tau_i, w_i \rangle_i = \sum_{n\in\mathbb{N}} \langle (e^i_n)^{\otimes 2}, w_i \rangle_i .$$

We recall that all function in two variables $(\xi_1, \xi_2) \in N_1 \times N_2$ is identified in an obvious manner with a single-variable function on the direct sum $N = N_1 \oplus N_2$, which is again a countably Hilbert nuclear space. Then, for $F \in C^2(N)$, (3.1) is written, for some $p, q \in \mathbb{N}$, in the following form:

$$\begin{aligned} F(\xi_1 + \eta_1, \xi_2 + \eta_2) &= F(\xi_1, \xi_2) + \sum_{i=1}^{2} \langle F'_i(\xi_1, \xi_2), \eta_i \rangle + \frac{1}{2} \sum_{i,j=1}^{2} \langle F''_{i,j}(\xi_1, \xi_2)\eta_i, \eta_j \rangle \\ &+ o(|\eta_1|_p^2 + |\eta_2|_q^2), \end{aligned} \tag{3.7}$$

where $F'_i(\xi_1, \xi_2) \in N'_i$, $F''_{i,j}(\xi_1, \xi_2) \in \mathcal{L}(N_i, N'_j)$ for $i, j = 1, 2$ and the error term satisfies

$$\lim_{t\to 0} \frac{o(t(|\eta_1|_p^2 + |\eta_2|_q^2))}{t^2} = 0.$$

For $i = 1, 2$, identifying $(N_i)_0$ with a subspace of $H = (N_1)_0 \oplus (N_2)_0$ in the canonical manner, we write e^1_n and e^2_n for $e^1_n \oplus 0$ and $0 \oplus e^2_n$, respectively, for all $n \in \mathbb{N}$.

Theorem 3.2 *For any test function $\varphi(x, y) = \sum_{n,m\in\mathbb{N}} \langle x^{\otimes n} \otimes y^{\otimes m}, \varphi_{n,m} \rangle$ in $\mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$, the Gross Laplacian is given by:*

$$\begin{aligned} \Delta_G \varphi(x, y) &:= trace_H D^2 \varphi(x, y) \\ &= \sum_{n,m\in\mathbb{N}} \langle x^{\otimes^n} \otimes y^{\otimes^m}, \langle (n + 2)(n + 1)\tau_1, \varphi_{n+2,m} \rangle\rangle \\ &+ \sum_{n,m\in\mathbb{N}} \langle x^{\otimes^n} \otimes y^{\otimes^m}, \langle (m + 2)(m + 1)\tau_2, \varphi_{n,m+2} \rangle\rangle. \end{aligned} \tag{3.8}$$

Moreover, the Gross Laplacian Δ_G is a convolution operator on $\mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2)$ into itself, i. e., there exists a distribution $\mathcal{T} = (\mathcal{T}_{n,m})_{n,m\in\mathbb{N}}$ such that

$$\Delta_G(\varphi) = \mathcal{T} * \varphi,\ \varphi \in \mathcal{F}_{(\theta_1,\theta_2)}(N'_1 \times N'_2), \tag{3.9}$$

where the Taylor expansion of the distribution $\mathcal{T}$ is given by

$$\mathcal{T}_{n,m} = \begin{cases} \tau_1,\ n = 2,\ m = 0, \\ \tau_2,\ n = 0,\ m = 2, \\ 0,\ otherwise. \end{cases} \tag{3.10}$$

Proof. Let $\varphi(x_1,x_2)=\sum\limits_{n,m\in\mathbb{N}}\langle x_1^{\otimes n}\otimes x_2^{\otimes m},\varphi_{n,m}\rangle$ be a test function in $\mathcal{F}_{(\theta_1,\theta_2)}(N_1'\times N_2')$. So, for all $(x_1,x_2),(y_1,y_2)\in N_1'\times N_2'$, the Taylor expansion of the function $\varphi(x_1+y_1,x_2+y_2)$ is given by

$$\begin{aligned}
\varphi(x_1+y_1,x_2+y_2) &= \sum_{n,m\in\mathbb{N}}\langle (x_1+y_1)^{\otimes^n}\otimes(x_2+y_2)^{\otimes^m},\varphi_{n,m}\rangle\\
&= \sum_{n,m,i,j\in\mathbb{N}}\begin{pmatrix} n+i\\ i\end{pmatrix}\begin{pmatrix} m+j\\ j\end{pmatrix}\langle x_1^{\otimes^n}\otimes x_2^{\otimes^m},\langle y_1^{\otimes i}\otimes y_2^{\otimes j},\varphi_{n+i,m+j}\rangle_{i,j}\rangle\\
&= \varphi(x_1,x_2)+\sum_{n,m\in\mathbb{N}}\langle x_1^{\otimes^n}\otimes x_2^{\otimes^m},\langle y_1,(n+1)\varphi_{n+1,m}\rangle_{1,0}+\langle y_2,(m+1)\varphi_{n,m+1}\rangle_{0,1}\rangle\\
&+\sum_{i,j=1}^{2}\sum_{n,m\in\mathbb{N}}\begin{pmatrix} n+i\\ i\end{pmatrix}\begin{pmatrix} m+j\\ j\end{pmatrix}\langle x_1^{\otimes^n}\otimes x_2^{\otimes^m},\langle y_1^{\otimes i}\otimes y_2^{\otimes j},\varphi_{n+i,m+j}\rangle_{i,j}\rangle+\varepsilon(y_1,y_2),
\end{aligned}\tag{3.11}$$

where for each $(x_1,x_2)\in N_1'\times N_2'$, ε is given by

$$\varepsilon(y_1,y_2)=\sum_{\substack{n,mi,j\in\mathbb{N}\\ i,j\geq 3}}\begin{pmatrix} n+i\\ i\end{pmatrix}\begin{pmatrix} m+j\\ j\end{pmatrix}\langle x_1^{\otimes^n}\otimes x_2^{\otimes^m},\langle y_1^{\otimes i}\otimes y_2^{\otimes j},\varphi_{n+i,m+j}\rangle_{i,j}\rangle.$$

It is easy to see that the function ε satisfy

$$\lim_{t\to 0}\frac{\varepsilon\left(t(y_1,y_2)\right)}{t}=0.$$

Identifying the equality (3.11) to the development (3.7), the second derivative of φ at (x_1,x_2) in the direction $(y_1,y_2)\in H$ is given by

$$D^2\varphi(x_1,x_2)=\sum_{i,j=1}^{2}\sum_{n,m\in\mathbb{N}}\begin{pmatrix} n+i\\ i\end{pmatrix}\begin{pmatrix} m+j\\ j\end{pmatrix}\langle x_1^{\otimes^n}\otimes x_2^{\otimes^m},\langle y_1^{\otimes i}\otimes y_2^{\otimes j},\varphi_{n+i,m+j}\rangle_{i,j}\rangle.$$

Using Lemma 3.1, we have for all $(x,y)\in N_1'\times N_2'$:

$$\begin{aligned}
trace_H D^2\varphi(x,y) &= \sum_{i,j\in\mathbb{N}}\langle D^2\varphi(x,y)(e_i^1,e_j^2),e_i^1\otimes e_j^2\rangle\\
&= \sum_{n,m,i,j\in\mathbb{N}}\langle x^{\otimes^n}\otimes y^{\otimes^m},(n+2)(n+1)\langle (e_i^1)^{\otimes 2},\varphi_{n+2,m}\rangle_{2,0}+(m+2)(m+1)\langle (e_j^2)^{\otimes 2},\varphi_{n,m+2}\rangle_{0,2}\rangle\\
&= \sum_{n,m\in\mathbb{N}}\langle x^{\otimes^n}\otimes y^{\otimes^m},\langle((n+2)(n+1)\tau_1,\varphi_{n+2,m}\rangle_{2,0}+(m+2)(m+1)\tau_2,\varphi_{n,m+2}\rangle_{0,2})\rangle.
\end{aligned}\tag{3.12}$$

In the other hand, using Lemma 2.2 for the distribution $\mathcal{T}\in\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1'\times N_2')$ where the Taylor expansion is given in (3.10) and the equality (3.12), we obtain

$$\begin{aligned}
\mathcal{T}*\varphi(x,y) &= \sum_{n,m\in\mathbb{N}}\langle x^{\otimes^n}\otimes y^{\otimes^m},\langle \mathcal{T}_{2,0},\varphi_{n+2,m}\rangle_{2,0}+\mathcal{T}_{0,2},\varphi_{n,m+2}\rangle_{0,2}\rangle\\
&= \sum_{n,m\in\mathbb{N}}\langle x^{\otimes^n}\otimes y^{\otimes^m},\langle((n+2)(n+1)\tau_1,\varphi_{n+2,m}\rangle_{2,0}+(m+2)(m+1)\tau_2),\varphi_{n,m+2}\rangle_{0,2}\rangle\\
&= \Delta_G\varphi(x,y),\ (x,y)\in N_1'\times N_2',\ \varphi\in\mathcal{F}_{(\theta_1,\theta_2)}(N_1'\times N_2').
\end{aligned}$$

Since the Gross Laplacian Δ_G is a convolution operator then

$$\Delta_G \in \mathcal{L}(\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2'), \mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')).$$

Remark 3.1 *Let $\varphi \in \mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$. Then, the Taylor expansion (3.8) of Δ_G, can be written as follows*

$$\Delta_G \varphi = \Delta_G^1 \varphi + \Delta_G^2 \varphi, \tag{3.13}$$

where for $i = 1, 2$, $\Delta_G^i \varphi$ is the Gross Laplacian with respect to the variable $\xi_i \in N_i'$ given by

$$\Delta_G^1 \varphi(x_1, x_2) = \sum_{n,m\in\mathbb{N}} \langle x_1^{\otimes^n} \otimes x_2^{\otimes^m}, \langle (n+2)(n+1)\tau_1, \varphi_{n+2,m}\rangle\rangle$$

respectively,

$$\Delta_G^2 \varphi(x_1, x_2) = \sum_{n,m\in\mathbb{N}} \langle x_1^{\otimes^n} \otimes x_2^{\otimes^m}, \langle (m+2)(m+1)\tau_1, \varphi_{n,m+2}\rangle\rangle.$$

Using the previous remark, it is easy to prove the following result.

Proposition 3.3 *For $\varphi = f \otimes g \in \mathcal{F}_{\theta_1}(N_1') \otimes \mathcal{F}_{\theta_2}(N_2') \subset \mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$, we obtain*

$$\Delta_G \varphi = \Delta_G(f \otimes g) = \Delta_G(f) \otimes g + f \otimes \Delta_G(g).$$

Therefore, on the subspace $\mathcal{F}_{\theta_1}(N_1') \otimes \mathcal{F}_{\theta_2}(N_2')$ of $\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$, the Gross Laplacian Δ_G is given by

$$\Delta_G = (\Delta_G)_{|\mathcal{F}_{\theta_1}(N_1')} \otimes I_2 + I_1 \otimes (\Delta_G)_{|\mathcal{F}_{\theta_2}(N_2')},$$

where for $i = 1, 2$, $(\Delta_G)_{|\mathcal{F}_{\theta_i}(N_i')}$ (respectively I_i) is the classical Gross Laplacian (respectively the identity operator) acting on $\mathcal{F}_{\theta_i}(N_i')$.

Then, we obtain the following corollary:

Corollary 3.4 *For all $(\xi_1, \xi_2) \in N_1 \times N_2$, we have*

$$\Delta_G(e_{(\xi_1,\xi_2)}) = \Delta_G(e_{\xi_1} \otimes e_{\xi_2}) = (\langle \xi_1, \xi_1\rangle_1 + \langle \xi_2, \xi_2\rangle_2) e_{(\xi_1,\xi_2)}. \tag{3.14}$$

Proof. To prove the corollary, it is sufficient to see that for $i = 1, 2$, we have:

$$\begin{aligned} (\Delta_G)_{|\mathcal{F}_{\theta_i}(N_i')}(e_{\xi_i})(x_i) &= \sum_{n\in\mathbb{N}} \langle x_i^{\otimes n}, \langle \tau_i, \xi_i^{\otimes n+2}\rangle_2\rangle \\ &= \langle \xi_i, \xi_i\rangle_i \sum_{n\in\mathbb{N}} \langle x_i^{\otimes n}, , \xi_i^{\otimes n}\rangle \\ &= \langle \xi_i, \xi_i\rangle_i e_{\xi_i}(x_i), \end{aligned}$$

for all $\xi_i \in N_i$ and $x_i \in N_i'$. □

For $i = 1, 2$, we assume that the Young function θ_i satisfies the following condition:

$$\limsup_{x\to\infty} \frac{\theta_i(x)}{x^2} < +\infty, \tag{3.15}$$

we obtain the following Gel'fand triple (see [5])

$$\mathcal{F}_{\theta_i}(N_i') \to L^2(X_i', \gamma_i) \to \mathcal{F}^*_{\theta_i}(N_i'), \tag{3.16}$$

$$\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2') \to L^2(X_1' \times X_2', \gamma_1 \otimes \gamma_2) \to \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2'), \tag{3.17}$$

where γ_i is the Gaussian measure on the real Fréchet nuclear space X_i' whose complexification is N_i defined via the Bochner-Minlos theorem [7] by its characteristic function:

$$\int_{X_i'} e^{i\langle x,\xi_i\rangle_i} d\gamma(x) = e^{-\frac{1}{2}|\xi_i|_0^2}, \xi_i \in X_i. \tag{3.18}$$

Using the Gel'fand triplet (3.17) and the fact that the Gross Laplacian is a convolution product, we extend the Gross Laplacian on $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$ as follows:

Definition 3.1 *The generalized Gross Laplacian acting on the distributions space $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$ is defined by*

$$\Delta_G(\Psi) = \mathcal{T} * \Psi, \ \Psi \in \mathcal{F}^*_{(\theta_1\theta_2)}(N_1' \times N_2'). \tag{3.19}$$

Proposition 3.5 *The Gross Laplacian Δ_G on $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$ coincide with the adjoint of the Gross Laplacian Δ_G^* on the test functions space $\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$.*

Proof. For any $\Phi \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$ and $\varphi \in \mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$, we have

$$\begin{aligned} \langle\langle \Delta_G \Phi, \varphi \rangle\rangle &= \langle\langle \mathcal{T} * \Phi, \varphi \rangle\rangle \\ &= [(\mathcal{T} * \Phi) * \varphi](0) = [\Phi * (\mathcal{T} * \varphi)](0) \\ &= \langle\langle \Phi, \mathcal{T} * \varphi \rangle\rangle = \langle\langle \Phi, \Delta_G(\varphi) \rangle\rangle \\ &= \langle\langle \Delta_G^*(\Phi), \varphi \rangle\rangle. \end{aligned}$$

4 Quantum Gross Laplacian

From the nuclearity of the space $\mathcal{F}_{\theta_1}(N_1')$, we have by Schwartz-Grothendieck Kernel Theorem the following isomorphisms

$$\mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2')) \simeq \mathcal{F}^*_{\theta_1}(N_1') \hat{\otimes} \mathcal{F}^*_{\theta_2}(N_2') \simeq \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2'). \tag{4.1}$$

So, for every $\Xi \in \mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2'))$, the associated kernel $\Xi^K \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$ is defined by

$$\langle\langle \Xi\varphi, \psi \rangle\rangle = \langle\langle \Xi^K, \varphi \otimes \psi \rangle\rangle, \ \forall \varphi, \psi \in \mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2'). \tag{4.2}$$

In the sequel, we will identify every operator Ξ with its kernel Ξ^K.

For every $\Xi \in \mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2'))$, the symbol is defined in the usual manner (see [12, 13]) by

$$\sigma(\Xi)(\xi_1, \xi_2) = \langle\langle \Xi e_{\xi_1}, e_{\xi_2} \rangle\rangle = \langle\langle \Xi^K, e_{(\xi_1,\xi_2)} \rangle\rangle = L(\Xi^K)(\xi_1, \xi_2) \tag{4.3}$$

where $\xi_1 \in N_1$, $\xi_2 \in N_2$. Then, every operator in $\mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2'))$ is uniquely determined by its symbol since the exponential vectors $e_{\xi_1} \otimes e_{\xi_2}$ span a dense subspace of $\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$. As direct application of Theorem 2.1 and the identification (4.2) by the kernel theorem, we obtain the following characterization of operators:

Theorem 4.1 *[3] The symbol map $\Xi \mapsto \sigma(\Xi)$ is a topological isomorphism:*

$$\mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2')) \quad \rightarrow \quad \mathcal{G}_{(\theta_1^*, \theta_2^*)}(N_1 \times N_2).$$

Let Ξ_1, $\Xi_2 \in \mathcal{L}(\mathcal{F}_\theta(N'), \mathcal{F}_\theta(N'))$, we define the convolution product of Ξ_1 and Ξ_2 denoted by $\Xi_1 * \Xi_2$ by

$$\sigma(\Xi_1 * \Xi_2) = \sigma(\Xi_1)\sigma(\Xi_2). \tag{4.4}$$

Using the topological isomorphisms:

$$\mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2')) \ni \Xi \mapsto \Xi^K \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2') \tag{4.5}$$

defined in (4.1) and the extended Gross Lapalcian given in (3.19), we can define the quantum Gross Laplacian as follows.

Definition 4.1 *The quantum Gross Laplacian Δ_G^Q is defined by:*

$$\Delta_G^Q(\Xi) = \mathcal{T} * \Xi^K, \; \Xi \in \mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2')), \tag{4.6}$$

where $\mathcal{T}$ is defined by (3.10).

Proposition 4.2 *For all $\Xi \in \mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2'))$, the quantum Gross Laplacian Δ_G^Q has the following analytic characterization:*

$$\sigma(\Delta_G^Q(\Xi))(\xi_1, \xi_2) = (\langle \xi_1, \xi_1 \rangle_1 + \langle \xi_2, \xi_2 \rangle_2)\sigma(\Xi)(\xi_1, \xi_2), \; (\xi_1, \xi_2) \in N_1 \times N_2. \tag{4.7}$$

Proof. Using the property (4.4) of the symbol of the convolution product of two operators and the definition (4.3), we have

$$\sigma(\Delta_G^Q(\Xi)) = L(\mathcal{L} * \Xi^K) = L(\mathcal{T})L(\Xi^K) = L(\mathcal{T})\sigma(\Xi), \; \Xi \in \mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2')).$$

In the other hand, using the Taylor expansion of $\mathcal{T}$, we obtain for all $(\xi_1, \xi_2) \in N_1' \times N_2'$

$$\begin{aligned} L(\mathcal{T})(\xi_1, \xi_2) &= \langle \mathcal{T}_{2,0}, \xi_1^{\otimes 2} \rangle_1 + \langle \mathcal{T}_{0,2}, \xi_2^{\otimes 2} \rangle_2 \\ &= \langle \tau_1, \xi_1^{\otimes 2} \rangle_1 + \langle \tau_2, \xi_2^{\otimes 2} \rangle_2 \\ &= \langle \xi_1, \xi_1 \rangle_1 + \langle \xi_2, \xi_2 \rangle_2. \end{aligned}$$

This proves the proposition.

5 Relation between classical and quantum Gross Laplacian

In this section, we consider the particular case where $\theta_1 = \theta_2 = \theta$ and $N = N_1 = N_2$. For simplicity, we use the same notation for the Gross Laplacian acting on $\mathcal{F}_\theta(N')$ and on $\mathcal{F}_{(\theta_1,\theta_2)}(N_1' \times N_2')$. It is well known (see [5]), that the pointwise multiplication yields a continuous bilinear map from $\mathcal{F}_\theta(N') \times \mathcal{F}_\theta(N')$ into $\mathcal{F}_\theta(N')$, i. e., for all $f, g \in \mathcal{F}_\theta(N')$, we have $fg \in \mathcal{F}_\theta(N')$. Let $\Phi \in \mathcal{F}^*_\theta(N')$ fixed and $\mathcal{M}_\Phi \in \mathcal{L}(\mathcal{F}_\theta(N'), \mathcal{F}^*_\theta(N'))$ be the multiplication operator by Φ defined by

$$\langle\!\langle \mathcal{M}_\Phi f, g \rangle\!\rangle = \langle\!\langle \Phi, fg \rangle\!\rangle, \; f, g \in \mathcal{F}_\theta(N'). \tag{5.1}$$

Let $e_0 = (1, 0, \ldots) \in \mathcal{F}_\theta(N')$ be the vacuum vector. Using the definition (5.1), we have the following connection between the multiplication operator and a distribution given by

$$\mathcal{M}_\Phi e_0 = \Phi.$$

Therefore, we obtain the following relation between classical and quantum Gross Laplacian.

Proposition 5.1 *For any $\Phi \in \mathcal{F}_\theta^*(N')$, we have:*

$$\left(\Delta_G^Q \mathcal{M}_\Phi\right) e_0 = \Delta_G \Phi. \tag{5.2}$$

Proof. Let $\Phi \in \mathcal{F}_\theta^*(N')$. Using the definitions of the Laplace transform (2.8) and the multiplication operator (5.1), we have

$$\begin{aligned}
\sigma(\mathcal{M}_\Phi)(\xi, \eta) &= \langle\!\langle \mathcal{M}_\Phi e_\xi, e_\eta \rangle\!\rangle \\
&= \langle\!\langle \Phi, e_{\xi+\eta} \rangle\!\rangle \\
&= \mathrm{L}(\Phi)(\xi + \eta),\ \xi, \eta \in N.
\end{aligned} \tag{5.3}$$

In the other hand, the Laplace transform of the classical Gross Lapalcian is given by

$$L(\Delta_G(\Phi))(\eta) = L(\mathcal{T} * \Phi)(\eta) = \langle \eta, \eta \rangle \widehat{\Phi}(\eta),\ \eta \in N.$$

By Proposition 4.2 and the equality (5.3), we obtain

$$\begin{aligned}
L(\Delta_G^Q(\mathcal{M}_\Phi)(e_0))(\eta) &= \langle\langle \Delta_G^Q(\mathcal{M}_\Phi)(e_0), e_\eta \rangle\rangle \\
&= \langle\langle \Delta_G^Q \mathcal{M}_\Phi, e_0 \otimes e_\eta \rangle\rangle \\
&= \sigma(\Delta_G^Q(\mathcal{M}_\Phi))(0, \eta) \\
&= \langle \eta, \eta \rangle \sigma(\mathcal{M}_\Phi))(0, \eta) \\
&= \langle \eta, \eta \rangle \widehat{\Phi}(\eta) \\
&= L(\Delta_G(\Phi))(\eta),\ \eta \in N.
\end{aligned}$$

Using the fact that the Laplace transform is an isomorphism, we have

$$\Delta_G^Q \mathcal{M}_\Phi e_0 = \Delta_G \Phi,\ \Phi \in \mathcal{F}_\theta^*(N').$$

6 Solution of linear quantum stochastic differential equations

Let $I \subset \mathbb{R}$ be an interval containing the origin. Consider a family $\{\Phi_t; t \in I\}$ of distributions in $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$ i. e., the function $t \mapsto \Phi_t$ is continuous from I into $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$. So, by using the isomorphism between $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$ and $\mathcal{G}_{(\theta_1^*,\theta_2^*)}(N_1 \times N_2)$ via the Laplace transform, the function $t \mapsto \widehat{\Phi}_t$ is continuous from I into $\mathcal{G}_{(\theta_1^*,\theta_2^*)}(N_1 \times N_2)$. Then, for each $t \in I$, the set $\{\widehat{\Phi}_s; s \in [0, t]\}$ is a compact subset of the generalized space $\mathcal{G}_{(\theta_1^*,\theta_2^*)}(N_1 \times N_2)$. This implies that it is bounded in $\mathcal{G}_{(\theta_1^*,\theta_2^*)}(N_1 \times N_2)$. Hence, there exist constants $p, q \in \mathbb{N}$, $a_1, a_2 > 0$ and $C_t > 0$ such that

$$|\widehat{\Phi}_t(\xi, \eta)| \leq C_t e^{\theta_1^*(a_1|\xi|_p)+\theta_2^*(a_2|\eta|_q)}|,\ (\xi, \eta) \in N_1 \times N_2.$$

By consequence, the function $(\xi,\eta) \mapsto \int_0^t \Phi_s(\xi,\eta)ds$ and belongs to the space $\mathcal{G}_{(\theta_1^*,\theta_2^*)}(N_1 \times N_2)$. This way we can define the integral $E_t = \int_0^t \Phi_s(\xi,\eta)ds$ as the unique element of $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$ satisfying

$$\left(\int_0^t \Phi_s(\xi,\eta)ds\right)^{\frown} = \int_0^t \widehat{\Phi}_s(\xi,\eta)ds,\ (\xi,\eta) \in N_1 \times N_2.$$

Moreover, for $t \in I$, the process E_t is differentiable in $\mathcal{F}_\theta(N' \times N')^*$ and satisfies the equation

$$\frac{\partial E_t}{\partial t} = \Phi_t.$$

Proposition 6.1 *For every distribution $\Phi \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2')$, the functional $e^{*\Phi}$ is defined by*

$$\widehat{e^{*\Phi}} = e^{\widehat{\Phi}} \tag{6.1}$$

belongs to $\mathcal{F}^*_{\left((e^{\theta_1^*})^*,(e^{\theta_2^*})^*\right)}(N_1' \times N_2')$.

Proof. The proof is similar to the one of Theorem 1 established in the paper [4]. □

Consider the following initial value problem:

$$\begin{cases} \frac{d\Xi(t)}{dt} = Z(t) * \Xi(t) + \Theta(t) \\ \Xi(0) = \Xi_0, \end{cases} \tag{6.2}$$

where $t \mapsto Z(t)$ and $t \mapsto \Theta(t)$ are continuous map defined on an interval I into $\mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2'))$ and $\Xi_0 \in \mathcal{L}(\mathcal{F}_{\theta_1}(N_1'), \mathcal{F}^*_{\theta_2}(N_2'))$.

Theorem 6.2 *The stochastic quantum differential equation (6.2) has a unique solution in $\mathcal{L}(\mathcal{F}_{\left(e^{\theta_1^*}\right)^*}(N_1'), \mathcal{F}^*_{\left(e^{\theta_2^*}\right)^*}(N_2'))$ given by*

$$\Xi(t) = \Xi_0 * e^{*\int_0^t Z(s)ds} + \int_0^t e^{*(\int_s^t Z(u)du)} * \Theta_s^K ds. \tag{6.3}$$

Proof. Applying the symbol map to the differential equation (6.2), we obtain an ordinary differential equation given by

$$\begin{cases} \frac{d\sigma(\Xi)(t)}{dt} = \sigma(Z)(t)\sigma(\Xi)(t) + \sigma(\Theta)(t) \\ \sigma(\Xi)(0) = \sigma(\Xi_0) \in \mathcal{F}^*_{(\theta_1,\theta_2)}(N_1' \times N_2'), \end{cases} \tag{6.4}$$

whose solution is given by

$$\sigma(\Xi)(t) = \sigma(\Xi_0)e^{\int_0^t \sigma(Z)(s)ds} + \int_0^t e^{\int_s^t \sigma(Z)(u)du}\sigma(\Theta)(s)ds.$$

Moreover, by Proposition 6.1, $\sigma(\Xi)(t)$ is an element of $\mathcal{F}^*_{\left((e^{\theta_1^*})^*,(e^{\theta_2^*})^*\right)}(N_1' \times N_2')$, for all $t \in I$. Using the property (4.4), the solution of the linear quantum stochastic differential equation is given by

$$\Xi(t) = \Xi_0 * e^{*\int_0^t Z(s)ds} + \int_0^t e^{*(\int_s^t Z(u)du)} * \Theta_s ds.$$

Now, by Theorem 4.1, we obtain that

$$\Xi(t) \in \mathcal{L}(\mathcal{F}_{\left(e^{\theta_1^*}\right)^*}(N_1'), \mathcal{F}^*_{\left(e^{\theta_2^*}\right)^*}(N_2')),$$

for all $t \in I$.

6.1 Solution of quantum Gross heat equation

In this section, we consider two Young functions θ_1 and θ_2, satisfying

$$\lim_{x\to\infty}\frac{\theta_i(x)}{x^2}<\infty,\ i=1,2.$$

Theorem 6.3 *The quantum Gross heat equation*

$$\begin{cases} \frac{d\Xi(t)}{dt}=\frac{1}{2}\Delta_G^Q\Xi(t)+\Theta(t) \\ \Xi(0)=\Xi_0, \end{cases} \tag{6.5}$$

has a unique solution in $\mathcal{L}(\mathcal{F}_{(e^{\theta_1^*})^*}(N_1'),\mathcal{F}^*_{(e^{\theta_2^*})^*}(N_2'))$ *given by*

$$\Xi(t)=\Xi_0*e^{*\frac{t}{2}\mathcal{T}}+\int_0^t e^{*\frac{t-s}{2}\mathcal{T}}*\Theta(s)ds. \tag{6.6}$$

Proof. For all $t\in I$, let $Z(t)=\mathcal{T}$, where $\mathcal{T}$ is the distribution given by (3.10). Then, the equation (6.9) becomes the quantum Gross heat equation given by

$$\begin{cases} \frac{d\Xi(t)}{dt}=\frac{1}{2}\mathcal{T}*\Xi(t)+\Theta(t)=\frac{1}{2}\Delta_G^Q\Xi(t)+\Theta(t) \\ \Xi(0)=\Xi_0. \end{cases} \tag{6.7}$$

Therefore, we apply Theorem 6.2 to get the unique solution in equation (6.9). We can further rewrite this solution in another way. For $t>0$, we define $\gamma_t=\gamma_1^t\otimes\gamma_2^t(.)=\gamma_1\otimes\gamma_2(./\sqrt{t})$, where $\gamma=\gamma_1\otimes\gamma_2$ is the standard Gaussian measure on the space X' defined in (3.18). It follows that the solution (6.9) can be expressed as

$$\Xi(t)=\Xi_0*\widetilde{\gamma}_t+\int_0^t\widetilde{\gamma}_{t-s}*\Theta(s)ds,$$

where $\widetilde{\gamma}$ is a positive distribution in $\mathcal{F}^*_{(\theta_1,\theta_2)}(N_1'\times N_2')$ given by

$$\langle\langle\widetilde{\gamma}_t,\varphi\rangle\rangle=\int_{X_1'\times X_2'}\varphi(x_1,x_2)d\gamma_t(x_1,x_2)=\int_{X_1'\times X_2'}\varphi(\sqrt{t}x_1,\sqrt{t}x_2)d\gamma_1(x_1)d\gamma_2(x_2),$$

where $\varphi\in\mathcal{F}_{(\theta_1,\theta_2)}(N_1'\times N_2')$. $\square$

It is easy to see from the definitions of the kernel operator (4.2) and the convolution product of the operators (4.4), that for all $\Xi_1,\Xi_2\in\mathcal{L}(\mathcal{F}_{\theta_1}(N_1'),\mathcal{F}^*_{\theta_2}(N_2'))$, we have

$$\Xi_1*\Xi_2=\Xi_1^K*\Xi_2^K.$$

Therefore combining Theorem 6.3 for the particular case where $\Theta=0$ and the Gel'fand triplet (3.17), we obtain the following result:

Corollary 6.4 *Let* $\Xi_0\in\mathcal{F}_{(\theta_1,\theta_2)}(N_1'\times N_2')$. *Then, the quantum Gross heat equation*

$$\begin{cases} \frac{d\Xi_t}{dt}=\frac{1}{2}\Delta_G^Q\Xi(t) \\ \Xi(0)=\Xi_0, \end{cases} \tag{6.8}$$

has a unique solution whose kernel is given by

$$\Xi_t^K(y_1,y_2)=\int_{X_1\times X_2}\Xi_0(y_1+\sqrt{t}x_1,y_2+\sqrt{t}x_2)d\gamma_1(x_1)d\gamma_2(x_2). \tag{6.9}$$

References

[1] **A. Barhoumi, H. H. Kuo and H. Ouerdiane**, *Generalized Gross heat equation with noises*, Soochow J. Math., V. 32, No. 1, (2006), 113-125.

[2] **D. M. Chung and U. C. Ji**, *Some Cauchy problems in white noise analysis and associated semigroups of operators*, Stochastic Anal. Appl., 17(1), (1999), 1-22.

[3] **M. Ben Chrouda and H. Ouerdiane**, *Algebra of operators on holomorphic functions and Applications*, Journal of Mathematical Physics, Analysis and Geometry, Vol. 5 (2002), 65-76.

[4] **M. Ben Chrouda, M. Eloued and H. Ouerdiane**, *Convolution calculus and applications to stochastic differential equations*, Soochow J. Math., 28(4), (2002), 375-388.

[5] **R. Gannoun, R. Hachaichi, H. Ouerdiane and A. Rezgui** , *Un théorème de dualité entre espaces de fonctions holomorphes à croissance exponentielle* , J. Func. Anal. vol. 171, no. 1, (2000), 1-14.

[6] **L. Gross**, *Potential theory on Hilbert space*, J. Funct. Anal. , 1, (1967), 123- 181.

[7] **T. Hida, H. H. Kuo, J. Potthoff, and L. Streit**, *White Noise. An Infinite Dimensional Calculus*, Kluwer Academic Publishers, Dordrecht, (1993).

[8] **T. Hida, N. Obata, and K. Saitô**, *Infinite-dimensional rotations and Laplacians in terms of white noise calculus*, Nagoya Math. J. , 128, (1992), 65-93.

[9] **U. C. Ji, N. Obata and H. Ouerdiane**, *Analytic Characterization of Generalized Fock Space Operators as Two-variable Entire Functions with Growth Condition*, World Scientific, vol. 5, No. 3, (2002), 395-407.

[10] **H. H. Kuo**, *On Laplacian operators of generalized Brownian functionals*, Lect. Notes in Math. 1203 (1986), 119-128.

[11] **H. H. Kuo**, *White noise distribution theory*, CRC Press, (1996).

[12] **N. Obata**, *White Noise Calculus and Fock Spaces*, Lecture Notes in Math. 1577 (1994).

[13] **H. Ouerdiane**, *Noyaux et symboles d'opérateurs sur des fonctionnelles analytiques gaussiennes*, Japan. J. Math. 21 (1995), 223-234.

[14] **H. Ouerdiane**, *Infinite dimensional entire functions and applications to stochastic differential equations*, Notices of th South African Math. Society, 35, No. 1 (2004), 23-45.

[15] **M. A. Piech**, *A fundamental solution of the parabolic equation on Hilbert space II. The semigroup property*, Trans. Amer. Math. Soc. 150, (1970), 257-286.